

**Inhibition of lipolysis: a novel explanation for the antipyretic and hypothermic actions
of Acetaminophen**

Shazma Bashir, Busayo Elegunde, Winston A. Morgan*.

The Medicines Research Group, School of Health, Sport and Bioscience, University of East
London, Romford Road, Stratford, London, E15 4 LZ. UK

* corresponding author; E-mail: w.a.morgan@uel.ac.uk; Tel: +44 (0) 208 223 4182;

Abstract

Background: Acetaminophen is both widely used to treat children with fever and is also responsible for thousands being hospitalised annually. Historically the antipyretic actions of acetaminophen was attributed to the inhibition of cyclooxygenase (COX-1/2) enzymes and more recently a novel COX-1 variant (COX-3) located in the brain. However, the evidence for acetaminophen-mediated COX inhibition remains contentious. This study assesses the impact of acetaminophen and other putative COX-3 inhibitors on the release of fatty acids during lipolysis as an alternative mechanism by which antipyretics could reduce body temperature during fever.

Experimental: 3T3-L1 adipocytes, primary brown adipocytes and isolated mitochondria were exposed to COX-3 inhibitors and lipolysis and mitochondrial electron transport chain function assessed.

Results: Acetaminophen, aminopyrine and antipyrine at 1-10 mM caused a significant decrease (up to 70%; $P < 0.01$, from control) in lipolysis within 1, 3 and 24 hours without affecting cell viability. The inhibition was observed regardless of where along its signalling pathway lipolysis was stimulated. All three compounds were found to significantly attenuate mitochondrial function by up to 30% for complex I and 40% for complex II ($P < 0.01$, from control).

Conclusions: These novel observations combined with the known limited inhibition of the COX enzymes by acetaminophen suggest both the antipyresis and hypothermia induced by acetaminophen and related compounds could be attributed to the direct inhibition of lipolysis and mitochondrial function, rather than cyclooxygenase inhibition centrally. Further these observations could provide new drug targets for reducing fever with the added bonus that fewer individuals being hospitalized by accidental acetaminophen overdose.

Keywords: acetaminophen, hypothermia, cyclooxygenase, lipolysis, mitochondria, electron transport chain.

Introduction

Acetaminophen (Paracetamol) is one of the oldest and most widely used treatments for pain, inflammation and Pyresis (fever) worldwide. With such a broad spectrum of activity many

1 questions remain about the exact mechanisms of action of acetaminophen. The analgesic
2 properties are thought to be due to actions centrally on opioid, serotonergic and cannabinoid
3 pathways (Sharma & Mehta, 2014) [1] . The anti-inflammatory and antipyretic actions of
4 acetaminophen has traditionally been attributed to inhibition of the cyclooxygenase (COX)
5 enzymes centrally (Hinz *et al.*, 2008) [2]. However there is now extensive evidence to show
6 that acetaminophen has only weak anti-inflammatory properties and this has been attributed
7 to its limited inhibition of the COX-1 and COX-2 enzymes (Chandrasekharan *et al.*, 2002;
8 Botting & Ayoub 2005[3] [4]). In mammals fever is associated with increased COX-2/PGE₂
9 induction and activity. With the lack of COX inhibition by acetaminophen another target for
10 the antipyretic actions of this widely used compound needed.

11 During the last 20 years, many studies have revealed that the administration of
12 acetaminophen (>100 mg/kg) to non-febrile rodents, for which there is no COX-2 induction
13 results in hypothermia. One explanation which initially gained significant support was that
14 acetaminophen is inhibiting a novel COX-1 variant (COX-3). It was proposed that the protein
15 (COX-3) which was suggested to be specifically inhibited by acetaminophen is constitutively
16 expressed in rodents to regulate body temperature (Chandrasekharan *et al.*, 2002; Ayoub *et al*
17 2004) [3] [5]. The COX-3 hypothesis was further supported by the observation that other
18 drugs such as aminopyrine and antipyrine which are also antipyretics, have been suggested to
19 be putative COX-3 inhibitors, also produce hypothermia in mice (Chandrasekharan *et al*,
20 2002; Ayoub *et al*, 2004) [3] [5]. However, the hypothermic actions of acetaminophen is also
21 observed in non-febrile animals such as the rat and in humans where the COX-3 protein
22 cannot theoretically be expressed (Foster *et al* 2016; Kis *et al*, 2005) [6] [7]. Consequently an
23 alternative hypothesis is needed to explain the acetaminophen-induced hypothermia observed
24 in non-febrile animals and this could provide an explanation for the antipyretic actions of
25 acetaminophen in humans.

26 In terms of thermoregulation, rodents (homeotherms) are capable of maintaining their core
27 body temperature (T_c) within a fairly constant range (36.0-37.5°C). However, due to their
28 large surface area to mass ratio, small mammals must increase their metabolic heat
29 production to regulate T_c when housed at temperatures below their thermoneutral zone
30 (Gordon, 2012) [8]. To maintain T_c, rodents rely on the stimulation of lipolysis, a catabolic
31 process resulting in the conversion of triglycerides (TG) stored in adipocytes to free fatty
32 acids (FA) and glycerol, the FA are then oxidised in the mitochondria to generate heat
33 (Gordon, 2012; Luo & Liu, 2016) [8] [9]. During periods of cold stress, in the case of

1 laboratory mice temperatures below 30°C, the process begins with the release of adrenergic
2 neurotransmitters from sympathetic fibres which binds to β_3 -receptors located on surface of
3 adipocytes leading to the activation of adenylate cyclase. This initiated an increase in cAMP
4 levels, the activation of Protein Kinase A (PKA) and the phosphorylation of the lipolytic
5 enzymes initiating lipolysis (Luo & Liu, 2016; Ueta *et al.*, 2012) [9] [10]. A similar process
6 of brown adipose tissue activation and lipolysis is also observed in humans following
7 prolonged cold exposure (Lesna' *et al.*, 1999) [11].
8
9
10
11
12

13 In most laboratory situations, mice are housed at around 20–24°C (Gordon, 1993, 2004) [12]
14 [13]. With their thermoneutral zone around 30°C they must respond to the mild cold stress by
15 increasing their basal metabolic rate to avoid hypothermia (Gordon, 1985, 1990, 1993) [14]
16 [15] [12]. It is therefore reasonable to hypothesize that animals housed under such conditions
17 may be more susceptible to hypothermia if exposed to compounds which even mildly
18 decrease the availability of FA and or compromise mitochondrial function.
19
20
21
22
23
24

25 To better understand the mechanisms by which acetaminophen and other related antipyretics
26 could either reduce fever or induce hypothermia in non-febrile animals. Studies were
27 undertaken to assess the impact of acetaminophen, aminopyrine and antipyrine on rate of
28 lipolysis in adipocytes and on the functioning of key complexes in the electron transport
29 chain of isolated mitochondrial.
30
31
32
33
34
35
36
37

38 **Methods and Materials**

43 **Chemicals**

44 All chemical were supplied by Sigma Chemical, Dorset, UK unless stated otherwise.
45
46
47
48
49
50

51 **Cell culture and differentiation of 3T3 L1 pre-adipocytes.**

52 The mouse 3T3-L1 pre-adipocyte cells (Public Health England, U.K.) were maintained under
53 standard culture conditions (37°C, 5% CO₂, 95% air) with medium containing DMEM
54 supplemented with 10% FBS, 2 mM glutamine, 10,000 units penicillin/ml, 10 mg
55 streptomycin/ml (5 ml/500 ml of medium) and 250 µg/ml amphotericin B (5 ml/500 ml of
56
57
58
59
60
61
62
63
64
65

1 medium). Cell were seeded at 10-20,000 cells/cm² and sub-cultured every 3 days at about 70-
2 80% confluent using 0.25% trypsin and 0.03% EDTA.
3

4 For differentiation the cells were allowed to become confluent then allowed to grow for four
5 additional days and this was treated as Day 0. At day 1, cells were placed in DMEM
6 differentiation medium with 10% fetal bovine serum, 1 µg/ml insulin, 0.5 mM
7 isobutylmethylxanthine (IBMX), 1 µM dexamethasone, 2 µM rosiglitazone for 48 hours. On
8 day 3, the differentiation medium was switched to DMEM containing 10% FBS and 1 µg/ml
9 insulin and changed after every 48 hours from this stage until fully differentiated 3T3-L1
10 adipocytes formation: Public Health England, U.K. (Zebisch *et al.*, 2012) [16].
11 Differentiation was confirmed by Oil Red O staining (Moreno-Aliaga & Matsumura, 1999)
12 [17]. Viability was assessed by the MTT assay (Van Munster, *et al.*, 1993) [18].
13
14
15
16
17
18
19
20
21

22 **Isolation of rat primary brown adipocytes**

23
24 Primary brown fat adipocytes was isolated from Wistar rats (200g) purchased from Envigo
25 UK. The animals were killed by cervical dislocation and brown adipose tissue from the
26 interscapular depots was dissected out and placed on a parafilm in a small volume of
27 Krebs/Ringer phosphate buffer. The cells were isolated as described by (Cannon &
28 Nedergaard, 2008) [19]. Once isolate the cells were incubated in a 24 well place and viability
29 and lipolysis determined.
30
31
32
33
34
35

36 **Determination of lipolysis assay in differentiated 3T3-L1 adipocytes and brown** 37 **adipocytes.3**

38
39 For all experiments the cells were incubated in serum-free medium for 2 hours (Schweiger *et*
40 *al.*, 2014) [20]. The cells were then exposed to different concentrations of the lipolysis
41 stimulants in medium containing 2% BSA (fatty acid free) and glycerol release was
42 measured after 1, 3 and 24 hours, for basal lipolysis no stimulant was added. In tests
43 involving the inhibitors (acetaminophen, aminopyrine and antipyrine) they were added 30
44 minutes before the stimulants. At the end of the incubation period the 25 µL of the incubation
45 medium would be taken for storage at -20°C for assay later. To determine the level of
46 glycerol the 25 µL of each sample was transferred to a new 96-well plate and 100 µL of
47 glycerol reagent was added and read at 540 nm (Sigma-Aldrich, U.K.). The concentration of
48 glycerol was determined from a calibration curve (Luo & Liu, 2016) [9].
49
50
51
52
53
54
55
56
57
58
59

60 **Isolation of mitochondria homogenate.**

1 Mitochondria was removed and isolated from Wistar rats (200g) as described by Sandoval-
2 Sandoval-Acuna, *et al.*, (2012) [21].
3
4
5
6

7 **Determination of Complex I activity**

8
9
10 The activity of NADH-ubiquinone oxidoreductase (complex 1) was determined by measuring
11 the reduction of 0.05mM of the electron acceptor 2,6-dinitrophenolindophenol (DCPIP) and
12 NADH (0.05mM) in the presence of either acetaminophen, aminopyrine or antipyrine at
13 600nm. Complex 1 activity was confirmed using complex 1 inhibitor rotenone (0.1mM). The
14 reaction mixture consisted of potassium phosphate buffer (35mM, pH 7.4) in the presence
15 and absence of KCN and antimycin A. The reaction was started by adding the mitochondria
16 homogenate (6.22mg/ml) and incubated for 10 minutes at room temperature (37°C).
17
18
19
20
21
22
23
24
25
26
27
28

29 **Determination of Complex II activity**

30
31 The activity of Succinate–Ubiquinone Oxidoreductase activity (Complex II) activity was
32 determined by measuring the reduction of DCPIP (0.05mM) in the presence of succinate (5-
33 100mM) and either acetaminophen, aminopyrine or antipyrine, or a range of antipyretic
34 agents at 600nm. The reaction mixture consisted of potassium phosphate buffer (35mM, pH
35 7.4) in the presence and absence of KCN (0.5mM) and antimycin A (0.5mM) over 5-100mM
36 of succinate. Complex II activity was confirmed using malonate (0.1mM).The reaction was
37 started by adding the mitochondria homogenate (6.22mg/ml) and incubated for 10 minutes at
38 room temperature (37°C).
39
40
41
42
43
44
45
46

47 The results were analysed using analysis of variance (ANOVA), followed by Dunnett's
48 Multiple Comparison Test. *P < 0.05, ** P<0.01, from control was considered statistically
49 significant.
50
51
52
53
54
55
56
57
58

59 **Results:**

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

The effect of Acetaminophen, Aminopyrine and Antipyrine on basal lipolysis in rat primary brown adipocytes.

Prior to the lipolysis studies, the cells were assessed for the impact of acetaminophen (PA), aminopyrine (AP) and antipyrine (AT) at concentrations up to 10 mM on cell viability. The concentrations of the antipyretics used in these studies were within the range normally associated with hypothermia *in vivo*: Fischer *et al.*, (1981) [22]; Orbach *et al.*, (2017) [23]. The compounds had no effect on the viability of primary adipocytes (24 hours) and up to 48 hours for 3T3-L1 cells (data not shown).

Glycerol release was assayed as an indicator of the extent of lipolysis Schweiger *et al.*, (2014) [20]. The primary brown fat adipocytes had a high level of basal glycerol release which did not significantly increase on stimulation. Treatment of primary brown adipocytes with acetaminophen at concentrations (1 mM and 10 mM) reduced basal glycerol release (10% and 19%) at 1 hour and (19% and 26%) at 24 hours (Figure 3.1). Similarly, aminopyrine (1 mM and 10 mM) attenuated glycerol release (22% and 17%) at 1 hour and (34% and 26%) at 24 hours (Figure 3.2). Antipyrine treatment resulted in a smaller but significant decrease in glycerol release (10% and 11%) at 1 hour and (21% and 19%) at 24 hours (Figure 3.3).

Effect of Acetaminophen on basal lipolysis in 3T3-L1 adipocytes

In unstimulated 3T3-L1 adipocytes incubated with acetaminophen at concentrations up to 10mM there was a significant decrease in glycerol release indicating a reduction in basal lipolysis similar to the primary brown adipocytes. This inhibitory effect can be seen from 1 hour with a 47% decrease in glycerol levels, followed by 57% and 52% at 3 and 24 hours respectively suggesting maximum lipolysis occurs within hours of exposure to acetaminophen (Figure 3.4).

Effect of Acetaminophen on Isoproterenol stimulated lipolysis in 3T3-L1 adipocytes

The effect of acetaminophen on lipolysis stimulated by isoproterenol a β -adrenergic agonist was then examined. Addition of isoproterenol elevated glycerol release from adipocytes as compared with that under basal conditions (Figure 3.5A). The glycerol release was both concentration and time dependent over 24 hours although the greater impact was with incubation time (Figure 3.5-3.6).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Addition of acetaminophen attenuated the glycerol release in a concentration dependent manner. Typically at the concentration of 0.001 μ M isoproterenol, 1mM of acetaminophen attenuated glycerol release by 41% whereas at 10mM acetaminophen the attenuation was 54%. A similar pattern was seen at 0.01 μ M and 0.1 μ M isoproterenol (Figure 3.5-6 B-D).

Effect of Acetaminophen on Forskolin stimulated lipolysis in 3T3-L1 adipocytes

In order to assess the post receptor impact of acetaminophen, the adenylate cyclase activator forskolin was used (Bezair *et al.*, 2009) [24]. Forskolin stimulated lipolysis in the 3T3-L1 cells (Figure 3.7). Prior exposure to acetaminophen (10 mM) attenuated the lipolysis induced by forskolin by 32%, 35% and 46% at 1, 3 and 24 hours in Figure 3.7 B,D,F).

Effect of acetaminophen on 8-Br-cAMP stimulated lipolysis in 3T3-L1 adipocytes

The cAMP analog; 8-Br-cAMP stimulated lipolysis in the 3T3-L1 cells (Figure 3.8). In the cells in the presence of acetaminophen (10 mM) glycerol release was significantly attenuated at 3 hours (45%) and still persisted (40%) at 24 hours (Figure 3.8B) confirming that the acetaminophen effect could also be located beyond the cAMP level.

Effect of Aminopyrine and Antipyrine on basal lipolysis in 3T3-L1 adipocytes

In addition to acetaminophen other putative COX-3 inhibitors have also been shown to induce hypothermia including aminopyrine and antipyrine. In this study both compounds attenuated basal lipolysis at 1 and 24 hours to an extent similar to acetaminophen (Figure 3.9).

The effect of Acetaminophen Aminopyrine and Antipyrine on mitochondrial activity.

Studies were also undertaken to determine the effect of acetaminophen, aminopyrine and antipyrine on the activity of complex I and complex II. At concentrations up to 10mM all the antipyretic compounds were found to inhibit complex I activity by decreasing the rate of rotenone sensitive NADH oxidation between 20-30% (Figure 3. 10). Similarly the

1 compounds also attenuated complex II activity as demonstrated by the inhibition of malonate
2 sensitive reduction DCPIP between 20-40% (Figure 3.11). The inhibition appears to be more
3 pronounced at higher succinate concentrations.
4
5
6
7

8 **Discussion:**

9
10 Acetaminophen is probably the most widely used over the counter antipyretic worldwide,
11 particularly for infants. Although it is very effective, there is still limited understanding about
12 how it lowers body temperature in humans and in febrile animal models. The situation is
13 further complicated because acetaminophen also causes hypothermia in non-febrile animals.
14 One consequence of our lack of understanding of the exact mechanisms of action of
15 acetaminophen is that every year thousands die or are hospitalised following acetaminophen
16 use. A better understanding would allow the development of safer alternatives
17
18
19
20
21
22
23
24
25
26
27

28 The focus of the present study is to understand how acetaminophen, and other putative COX-
29 3 inhibitors; aminopyrine and antipyrine could disrupt key cellular processes which
30 determine heat generation, with a view to gaining a novel insight into the mechanism of
31 action of antipyretics generally.
32
33
34
35

36 For homeotherms such as laboratory rodents housed at temperatures below their
37 thermoneutral zone there is a constant need to increase heat generation to avoid hypothermia.
38 When homeotherms detect the external temperature is below their thermoneutral zone, a
39 number of key responses are observed. One of the most important starts with signals from the
40 preoptic area of anterior hypothalamus (POA) which stimulates peripheral effectors leading
41 to the increase in lipolysis augmenting the amount of free fatty acids available for
42 mitochondrial metabolism and heat generation. Given the lack of agreement about the
43 effectiveness of acetaminophen at inhibiting cyclooxygenases centrally and peripherally, it is
44 reasonable to examine the impact of this key antipyretic at other points along the mammalian
45 heat generation pathway starting with lipolysis.
46
47
48
49
50
51
52
53
54

55 Lipolysis is a critical metabolic function for adipocytes and is the key process that provides
56 fatty acids for energy generation and to activate uncoupling protein 1, both are important for
57 thermogenesis (Ohlson *et al.*, 2004) [25]. Conversion of stored triglycerides into free fatty
58
59
60
61
62
63
64
65

1 acids (lipolysis) is stimulated by β -adrenergic signalling on the cell surface of adipocytes.
2 This leads to the activation of adenylate cyclase, increased levels of cAMP, the activation of
3 PKA and the phosphorylation and activation of lipolytic enzymes including *hormone*
4 *sensitive lipase*, *adipose triglyceride lipase*, and perilipin (Chrysovergis *et al.*, 2014;
5 McKnight *et al.*, 1998; MacPherson & Peters, 2015; Schweiger *et al.*, 2006; Heeren &
6 Műnzberg 2013; Marcelin & Chua, 2010) [26][27] [28] [29] [30] [31]. An essential
7 requirement for all these process, from cell surface receptor signalling pathway, through to
8 lipolysis and oxidation of the fatty acids, is the availability of ATP and functioning
9 mitochondria. Any disruption or inhibition of mitochondrial function could potentially
10 attenuate lipolysis with the consequent impact on heat generation leading to hypothermia.
11
12
13
14
15
16
17
18

19 In this study prior to the lipolysis studies, the 3T3-L1 adipocyte cells were assessed for any
20 direct impact of the antipyretics on viability. The concentrations selected (up to 10 mM) were
21 similar to or lower than the plasma concentrations known to induce hypothermia in animals
22 models and more importantly within the plasma concentrations required for antipyresis or
23 hypothermia in humans (Fischer *et al.*, 1981; Orbach *et al.*, 2017) [22] [23]. At these
24 concentrations over 24 hours, there was no loss of cell viability indicating that the effects
25 observed could not be attributed to the loss of adipocytes.
26
27
28
29
30
31

32 To examine whether the acetaminophen, aminopyrine and antipyrine could have a direct
33 impact on lipolysis, the extent of glycerol was assessed. This is the most frequently used *in*
34 *vitro* model and method to study lipolysis. To ensure detectable levels of lipolysis the
35 adipocytes were incubated in a low glucose medium which forces the cells to metabolise fats
36 for energy. Under basal conditions and without the presence of an adrenergic agonist,
37 acetaminophen at 10 mM was found to significantly reduce glycerol release as early as 1
38 hour. The fact that acetaminophen inhibited basal lipolysis suggested the effect could also be
39 at a post receptor site or totally independent of adrenergic stimulation.
40
41
42
43
44
45
46
47

48 To further investigate the impact of acetaminophen on the catecholamine receptor pathway,
49 the β -agonist isoproterenol was selected to stimulate lipolysis. Adrenergic stimulated
50 lipolysis is more likely to be observed in animals under cold stress such as rodents housed at
51 22°C. In these studies acetaminophen also inhibited isoproterenol induced lipolysis. This
52 observation prompted the question as to where along the adrenergic pathway acetaminophen
53 was acting. The attenuation of the basal lipolysis has already confirmed that at least some of
54 the actions acetaminophen is independent of adrenergic receptor binding. However, to
55
56
57
58
59
60
61
62
63
64
65

1 investigate whether acetaminophen was affecting the binding to the β -receptor, forskolin was
2 used. A direct activator of adenylate cyclase, forskolin was able to stimulate lipolysis and
3 which was also attenuated by acetaminophen (Bezair *et al.*, 2009) [24]. This observation
4 suggests acetaminophen inhibition is less likely to be at the β -receptor as forskolin work
5 downstream of receptor binding.
6
7

8
9
10 The next step was to investigate the effect of acetaminophen beyond the adenylate cyclase
11 stage of the signalling pathway. To determine the impact of acetaminophen post adenylate
12 cyclase, the cAMP analog 8-Br-cAMP was employed. This compound is able to enhance the
13 activity of protein kinase A thereby phosphorylating the lipolytic enzymes without activation
14 of adenylate cyclase. The addition of 8-Br-cAMP induced an increase in lipolysis and this
15 was attenuated by acetaminophen, confirming that the effect of must be distal to the
16 production of cAMP. These observation confirm that the impact of acetaminophen was not
17 on the adrenergic signalling pathway as acetaminophen inhibition was observed regardless of
18 where in the signally pathway lipolysis was stimulated.
19
20
21
22
23
24
25

26
27 In addition to acetaminophen, the effect of aminopyrine and antipyrine were also examined as
28 these compounds are known to cause hypothermia *in vivo* in mice and were reported to be
29 COX-3 inhibitors *in vitro* (Ayoub *et al* 2004; Chandrasekharan *et al*, 2002) [5] [3].
30 Antipyrine (phenazone) exhibits analgesic, antipyretic and antirheumatic activity.
31 Aminopyrine (aminophenazone) is thought to possess greater antipyretic and analgesic
32 activity, marked anti-inflammatory property but more toxic than antipyrine (Volz & Kellner,
33 1980) [32]. However like acetaminophen, little is understood about the impact of these
34 compounds on lipolysis and how this relates to thermogenesis. In the present study, both
35 aminopyrine and antipyrine were effective at significantly attenuating glycerol release in a
36 manner similar to acetaminophen. This is a novel observation and like acetaminophen could
37 be linked to antipyresis.
38
39
40
41
42
43
44
45
46
47

48 Although the differentiated 3T3-L1 cells are the most widely used *in vitro* model for studying
49 lipolysis and fat metabolism, investigations were also undertaken using freshly isolated rat
50 primary brown adipocytes. As primary cells they should provide a more realistic picture of
51 the effect of the test compound on lipolysis *in vivo*. The primary brown adipocytes cells
52 showed high levels of basal lipolysis which may reflect the fact that they had recently been
53 removed from animals that have acclimatized to living at ambient temperatures which are
54 below their thermoneutral zone and so require constant heat generation. The high level of
55
56
57
58
59
60
61
62
63
64
65

1 basal lipolysis made it difficult to increase the level of lipolysis. However the compounds
2 (acetaminophen, aminopyrine and antipyrine) inhibited lipolysis in the primary cells in a
3 manner similar to the 3T3-L1 cells. Given the surface area to mass ratio of small mammals
4 such as rodents any slight reduction heat generation would affect Tc. Consequently, when
5 such animals are housed at temperatures below their thermoneutral zone, any reduction in
6 lipolysis would compromise their ability to thermo-regulate. This may explain the
7 hypothermia observed when these animals are exposed to acetaminophen, aminopyrine and
8 antipyrine.
9

10
11
12
13
14
15 Although this is the first time acetaminophen, aminopyrine and antipyrine have been shown
16 to inhibit lipolysis they are not the only antipyretic compounds which have been shown to
17 inhibit lipolysis. Indomethacin has been shown to inhibit enzymes involved in basal and
18 stimulated lipolysis in the kidney (Erman *et al.*, 1980) [33]. Other NSAIDs (aspirin,
19 naproxen, nimesulide, and piroxicam) are known to activate the NADPH oxidase (NOX)
20 isoform (NOX4) in adipocytes to produce hydrogen peroxide (H₂O₂), which impairs cAMP-
21 dependent PKA-II activation, thus inhibiting isoproterenol activated lipolysis. H₂O₂
22 signalling is a novel COX independent effect of NSAIDs in adipocytes and may play a role in
23 antipyresis (Vázquez-Meza *et al.*, 2013) [34].
24
25
26
27
28
29
30
31

32 Although lipolysis is a key step in the thermogenesis process, acetaminophen and other
33 hypothermic compounds could also work at sites downstream to the release of fatty acids
34 from triglycerides, for example by directly inhibiting mitochondrial function. Given the lack
35 of inhibition of adrenergic signalling pathway in this study, further studies were conducted
36 using isolated mitochondria. As previously stated, energy generation for thermoregulation
37 require functioning mitochondria. In the present study all three compounds; acetaminophen,
38 aminopyrine and antipyrine were shown to attenuated mitochondrial activity at complex I and
39 complex II. Without fully functioning mitochondria both lipolysis and fatty acid oxidation
40 will be compromised, and this would directly lead to hypothermia in small mammals such as
41 rodents.
42
43
44
45
46
47
48
49
50

51 It has always been difficult to reconcile the pharmacological actions of acetaminophen
52 particularly the antipyretic effects with the weak inhibition of cyclooxygenase enzymes. The
53 novel observations in this study provide a logical and plausible explanation as to how
54 acetaminophen could cause both hypothermia and antipyresis distinct from any impact on
55 cyclooxygenase activity. Even at the levels of inhibition reported in this study this would
56
57
58
59
60
61
62
63
64
65

1 negatively impact on how small mammals thermoregulate at environmental temperatures
2 below their thermoneutral zone. These observations could explain the hypothermia widely
3 reported when some antipyretic compounds are administered to rodents at concentrations
4 normally required for antipyresis. Similarly during fever hyperthermia is driven by increased
5 metabolism including increased lipolysis and mitochondrial function. The attenuating effect
6 on lipolysis and mitochondrial function of these antipyretic agents observed in this study
7 could provide an alternative peripheral mechanism of the action for acetaminophen and many
8 other compounds which is independent of cyclooxygenase inhibition both centrally and
9 peripherally.
10
11
12
13
14
15
16
17
18

19 **Competing interests:** The authors declare that they have no competing interests.
20
21

22 **Funding:** All funding for all the authors came directly from the University of East London.
23
24
25

26 **Authors' contributions:** WM, and SB participated in the overall planning of experiments,
27 research design, data analysis & interpretation, and manuscript preparation. BE performed the
28 experiments on isolated mitochondria.
29
30
31
32
33

34 **DECLARATION OF TRANSPARENCY AND SCIENTIFIC RIGOUR**

35 This Declaration acknowledges that this paper adheres to the principles for transparent
36 reporting and scientific rigour of preclinical research as stated in the *BJP* guidelines for
37 [Design & Analysis](#), and [Immunoblotting and Immunochemistry](#), and as recommended by
38 funding agencies, publishers, and other organisations engaged with supporting research.
39
40
41
42
43
44
45
46
47
48

49 **References**

50
51
52
53
54
55

- 56 [1] Sharma, C.V., and Mehta, V., (2014). Acetaminophen: mechanisms and updates
57 Continuing Education in Anaesthesia, Critical. Care & Pain, 14 (4): 153-158.
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [2] Hinz, B., Cheremina, O., Brune, K., (2008). Acetaminophen (acetaminophen) is a selective cyclooxygenase-2 inhibitor in man. *FASEB J*, 22: 383-390.
- [3] Chandrasekharan, N.V., Dai, H., Roos, K.L., Evanson, N.K., Tomsik, J., Elton, T.S., and Simmons, D.L., (2002). COX-3, a cyclooxygenase-1 variant inhibited by acetaminophen and other analgesic/antipyretic drugs: cloning, structure, and expression. *Proc Natl Acad Sci U S A*, 99: 13926-13931.
- [4] Botting, R., and Ayoub, S.S., (2005). COX-3 and the mechanism of action of acetaminophen/acetaminophen. *Prostaglandins Leukot Essent Fatty Acids*, 72(2): 85-7.
- [5] Ayoub, S.S., Botting, R.M., Goorha, S., Colville-Nash, P.R., Willoughby, D.A., and Ballou L.R. (2004). Acetaminophen-induced hypothermia in mice is mediated by a prostaglandin endoperoxide synthase 1 gene-derived protein. *Proc Natl Acad Sci U S A*, 101(30): 11165-11169.
- [6] Foster, J., Mauger, A., Thomasson, K., White S., and Taylor L. (2016). Effect of Acetaminophen Ingestion on Thermoregulation of Normothermic, Non-febrile Hum. *Front Pharmacol*, 7: 54.
- [7] Kis, B., Snipes, J.A., and Busija, D.W., (2005). Acetaminophen and the cyclooxygenase-3 puzzle: sorting out facts, fictions, and uncertainties. *J. Pharmacol. Exp. Ther.* 315(1): 1-7.
- [8] Gordon, C.J., (2012). Thermal physiology of laboratory mice: Defining thermoneutrality. *Journal of Thermal Biology*, 37(8): 654-685.
- [9] Luo, L., and Liu M. (2016). Adipose tissue in control of metabolism. *J. Endocrinol.* 231(3): R77-R99.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [10] Ueta, C.B., Fernandes, G.W., Capelo, L.P., Fonseca, T.L., Maculan, F.D., Gouveia, C.H., Brum, P.C., Christoffolete, M.A., Aoki, M.S., Lancellotti, C.L., Kim, B., Bianco, A.C., and Ribeiro, M.O., (2012). β_1 Adrenergic receptor is key to cold- and diet-induced thermogenesis in mice. *J. of Endocrinol.* 214: 359–365.
- [11] Lesna', I., Vybi'ral, S., Jansky, L., Zeman, V., (1999). Human nonshivering thermogenesis. *J. Therm. Biol.* 24: 63–69.
- [12] Gordon, C.J., (1993). *Temperature regulation in laboratory rodents*. New York: Cambridge University Press. Xii, 276.
- [13] Gordon, C.J. (2004). Effect of cage bedding on temperature regulation and metabolism of group-housed female mice. *Comparative Medicine*, 54: 51-56.
- [14] Gordon, C.J., (1985). Relationship between autonomic and behavioral thermoregulation in the mouse. *Physiol. Behav*, 34: 687-690.
- [15] Gordon C.J. (1990). Thermal biology of the laboratory rat. *Physiol. Behav.* 47, 963-991.
- [16] Zebisch K., Voigt V., Wabitsch M., Brandsch M. (2012). Protocol for effective differentiation of 3T3-L1 cells to adipocytes. *Anal. Biochem.* 425(1): 88-90.
- [17] Moreno-Aliaga, M,J., Matsumura, F., (1999). Endrin inhibits adipocyte differentiation by selectively altering expression pattern of CCAAT/enhancer binding protein-alpha in 3T3-L1 cells. *Mol. Pharmacol.* 56: 91–101.
- [18] Van Munster, I.P., Tangerman, A., De Haan, A.F.J., and Nagengast, F.M., (1993). A new method for the determination of cytotoxicity of bile acids and aqueous phase of stool: the effect of calcium. *Euro. J. Clin. Invest.* 23: 773–777.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [19] Cannon B, Nedergaard J. (2008). Studies of thermogenesis and mitochondrial function in adipose tissues. *Methods Mol Biol.* 456:109-21.
- [20] Schweiger, M., Eichmann, T.O., Taschler, U., Zimmermann, R., Zechner, R., and Lass, A. (2014). Measurement of Lipolysis. *Meth. Enzymol.* 538: 171–193.
- [21] Sandoval-Acuna, C., Lopez-Alarcon, C., Aliaga, M.E., and Speisky, H., (2012). Inhibition of mitochondrial complex I by various non-steroidal anti-inflammatory drugs and its protection by quercetin via a coenzyme Q-like action. *Chem. Biol. Interact.* 199: 18-28.
- [22] Fischer, L.J., Green, M.D., and Harman, A.W., (1981). Levels of acetaminophen and its metabolites in mouse tissues after a toxic dose. *Journal of Pharmacology and Experimental Therapeutics*, 219: 281–286.
- [23] Orbach, S.M., Cassin, M.E., Ehrich, M.F., and Rajagopalan, P., (2017). Investigating acetaminophen hepatotoxicity in multi-cellular organotypic liver models. *Toxicol. In Vitr.* 42: 10-20.
- [24] Bezaire, V., Mairal, A., Ribet, C., Lefort, C., Girousse, A., Jocken, J., Laurencikiene, J., Anesia, R., Rodriguez, A., Ryden, M., Stenson, B.M., Dani C., Ailhaud, G., Peter Arner, P., Langin, D., (2009). Contribution of Adipose Triglyceride Lipase and Hormone-sensitive Lipase to Lipolysis in hMADS Adipocytes. *The Journal of Biological Chemistry*, 284 (27): 18282–18291.
- [25] Ohlson, K.B., Shabalina, I.G., Lennström, K., Backlund, E.C., Mohell, N., Bronnikov, G.E., Lindahl, S.G., Cannon, B., and Nedergaard J. (2004). Inhibitory effects of halothane on the thermogenic pathway in brown adipocytes: localization to adenylyl cyclase and mitochondrial fatty acid oxidation. *Biochem. Pharmacol.* 68(3): 463-477.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [26] Chrysovergis, K., Wang, X., Kosak, J., Lee, S-H., Kim, J.S., Foley, J.F., Travlos, G., Singh, S., Baek, S.J., and Eling, T.E., (2014). NAG-1/GDF15 prevents obesity by increasing thermogenesis, lipolysis and oxidative metabolism. *Int J Obes (Lond)*, 38(12): 1555–1564.
- [27] McKnight, G.S., Cummings, D.E., Amieux, P.S., Sikorski, M.A., Brandon, E.P., Planas, J.V., Motamed, K., and Idzerda, R.L. (1998). Cyclic AMP, PKA, and the physiological regulation of adiposity. *Recent. Prog. Horm. Res.* 53: 139–59.
- [28] MacPherson, R.E., and Peters, S.J. (2015). Piecing together the puzzle of perilipin proteins and skeletal muscle lipolysis. *Appl. Physiol. Nutr. Metab.* 40(7): 641-51.
- [29] Schweiger, M., Schreiber, R., Haemmerle, G., Lass, A., Fledelius, C., Jacobsen, P., Tornqvist, H., Zechner, R., and Zimmermann, R. (2006). Adipose triglyceride lipase and hormone-sensitive lipase are the major enzymes in adipose tissue triacylglycerol catabolism. *J. Biol. Chem.* 281: 40236–41.
- [30] Heeren, J., Münzberg, H., (2013). Novel aspects of brown adipose tissue biology. *Endocrinol. Metab. Clin.* 42: 89–107.
- [31] Marcelin, G., and Chua, S., (2010). Contributions of adipocyte lipid metabolism to body fat content and implications for the treatment of obesity. *Curr. Opin. Pharmacol.* 10: 588–593.
- [32] Volz, M., Kellner, H.M., (1980). Kinetics and metabolism of pyrazolones (propyphenazone, aminopyrine and dipyrone). *Br. J. Clin. Pharmacol.* 10 Suppl 2: 299S-308S.
- [33] Erman, A., Schwartzman, M., and Raz, A., (1980). Indomethacin but not aspirin inhibits basal and stimulated lipolysis in rabbit kidney. *Prostaglandins*, 20(4): 689-702.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

[34] Vázquez-Meza, H., de Piña, M.Z., Pardo, J.P., Riveros-Rosas, H., Villalobos-Molina, R., and Piña, E., (2013). Non-steroidal anti-inflammatory drugs activate NADPH oxidase in adipocytes and raise the H₂O₂ pool to prevent cAMP-stimulated Protein Kinase A activation and inhibit lipolysis. BMC Biochem.14: 13.

Figure Legends

Figure 3.1 Effect of acetaminophen on basal lipolysis in rat primary brown adipocytes at 1 and 24 hours: Brown adipocytes were treated with different concentrations of acetaminophen and glycerol release was measured at 1(A) and 24(B) hours. Data are representative of n=4 replicates expressed as means ± Standard deviations (**P < 0.01 from control).

Figure 3.2 Effect of aminopyrine on basal lipolysis in rat primary brown adipocytes at 1 and 24 hours: Brown adipocytes were treated with different concentrations of aminopyrine and glycerol release was measured at 1(A) and 24(B) hours. Data are representative of n=4 replicates expressed as means ± Standard deviations (**P < 0.01 from control).

Figure 3.3 Effect of antipyrine on basal lipolysis in rat primary brown adipocytes at 1 and 24 hours: Brown adipocytes were treated with different concentrations of antipyrine and glycerol release was measured at 1(A) and 24(B) hours. Data are representative of n=4 replicates expressed as means ± Standard deviations (**P < 0.01 from control).

Figure 3.4 Effect of acetaminophen on basal lipolysis in 3T3-L1 adipocytes at 1, 3 and 24 hours: Differentiated 3T3-L1 adipocytes were treated with serum free DMEM for 2 hours and were then treated with different concentrations of acetaminophen (PA). For basal

1 lipolysis, cells were treated at the same time with an appropriate volume of vehicle. Lipolysis
2 was determined by measuring glycerol released into the culture media at 1(A), 3(B) and
3 24(C) hours. Data are representative of n=4 replicates expressed as means \pm Standard
4 deviations (*P < 0.05, **P < 0.01 from control).
5
6
7
8
9

10 **Figure 3.5 Effect of acetaminophen on isoproterenol stimulated lipolysis in 3T3-L1**
11 **adipocytes at 1 hour:** Differentiated 3T3-L1 adipocytes were treated with serum free DMEM
12 for 2 hours. Cells were either treated with different concentrations of isoproterenol (Iso)
13 alone (A) or treated with different concentrations of acetaminophen and isoproterenol added
14 simultaneously (B-D). Lipolysis was determined by measuring glycerol released into the
15 culture media at 1 hour. Data are representative of n=4 replicates expressed as means \pm
16 Standard deviations (*P < 0.05, **P < 0.01 from control).
17
18
19
20
21
22
23
24
25
26

27 **Figure 3.6 Effect of acetaminophen on isoproterenol stimulated lipolysis in 3T3-L1**
28 **adipocytes after 24 hours:** Differentiated 3T3-L1 adipocytes were treated with serum free
29 DMEM for 2 hours. Cells were either treated with different concentrations of isoproterenol
30 alone (A) or treated with different concentrations of acetaminophen and isoproterenol added
31 simultaneously (B-D). Lipolysis was determined by measuring glycerol released into the
32 culture media at 24 hours. Data are representative of n=4 replicates expressed as means \pm
33 Standard deviations (*P < 0.05, **P < 0.01 from control).
34
35
36
37
38
39
40
41
42

43 **Figure 3.7 Effect of acetaminophen on forskolin stimulated lipolysis in 3T3-L1 adipocytes**
44 **at 1(A), 3(B) and 24(C) hours:** Differentiated 3T3-L1 adipocytes were treated with serum
45 free DMEM for 2 hours. Cells were either treated with forskolin (Fsk) alone (A) or
46 preincubated with different concentrations of acetaminophen for 30 minutes and then
47 forskolin added (B). Lipolysis was determined by measuring glycerol released into the
48 culture media at 1(A,B), 3(C,D) and 24(E,F) hours. Data are representative of n=4
49 replicates expressed as means \pm Standard deviations (*P < 0.05, **P < 0.01 from control).
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 3.8 Effect of acetaminophen on 8-Br-cAMP stimulated lipolysis in 3T3-L1 adipocytes at 3 and 24 hours: Differentiated 3T3-L1 adipocytes were treated with serum free DMEM for 2 hours. Cells were either treated with 8-Br-cAMP alone (A) or preincubated with different concentrations of acetaminophen for 30 minutes and then 8-Br-cAMP added (C). Lipolysis was determined by measuring glycerol released into the culture media at 3(A,C) and 24(B,D) hours. Data are representative of n=4 replicates expressed as means \pm Standard deviations (*P < 0.05, **P < 0.01 from control).

Figure 3.9 Effect of aminopyrine and antipyrine on basal lipolysis in 3T3-L1 adipocytes at 1, 3 and 24 hours: Differentiated 3T3-L1 adipocytes were treated with serum free DMEM for 2 hours and were then treated with aminopyrine (AM) or antipyrine (AT). Lipolysis was determined by measuring glycerol released into the culture media at 1(A), 3(B) and 24(C) hours. For basal lipolysis, cells were treated at the same time with an appropriate volume of vehicle. Data are representative of n=4 replicates expressed as means \pm Standard deviations (**P < 0.01 from control).

Fig 3. 10 The effect of antipyretic compounds on mitochondrial complex I activity.

Mitochondria were incubated with 5mM of either acetaminophen (Para) Antipyrine (AT) and Aminopyrine (AM) for 10 mins in the presence of NADH (0.05mM). The reduction of DCPIP at 600nm was monitored for 3 minutes. Values represents the mean \pm SD (N= 7). The significance of difference was determined using a one way ANOVA followed by a Dunnett's multiple of comparison post hoc test. **p< 0.001 was statistically significant compared to the control.


Fig 3.11 The effect of acetaminophen, aminopyrine and antipyrine on complex II activity in rat liver mitochondria. A concentration of 5mM of the test compounds was added to a reaction mixture containing DCPIP and mitochondria suspension. Varying concentration of succinate (20mM-100mM) was used and absorbance at 600nm was monitored for 3mins. Values are expressed as the mean \pm SEM (N= 16). *p<0.05 **p<0.01 and ***p <0.001

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

compared to respective control (succinate). AM=Aminopyrine, AT= Antipyrine and PARA = Acetaminophen.

Figure 3.1

A.



B.





Figure 3.2

A.



B.




Figure 3.3






Figure 3.4

A.



B.



C.




Figure 3.5




Figure 3.6




Figure 3.7





Figure 3.8

A.



B.




Figure 3.9




Figure 3.10





Figure 3.11



Highlights

- In terms of antipyresis and hypothermia, the peripheral actions of acetaminophen and related compounds are more important than inhibition of cyclooxygenases centrally.
- The antipyresis and hypothermic effects of acetaminophen may be due to the inhibition of lipolysis.
- The inhibition of mitochondrial electron transport chain by acetaminophen may explain both the antipyresis and hypothermic effects.