

University of East London Institutional Repository: <http://roar.uel.ac.uk>

This paper is made available online in accordance with publisher policies. Please scroll down to view the document itself. Please refer to the repository record for this item and our policy information available from the repository home page for further information.

To see the final version of this paper please visit the publisher's website. Access to the published version may require a subscription.

Author(s): Sholl, Catherine; Korkie, Juan; Harper, David J.

Article title: 'Challenging teenagers' ideas about people with mental health problems.

Year of publication: 2010

Citation: Sholl, C., Korkie, J. & Harper, D.J. (2010) 'Challenging teenagers' ideas about people with mental health problems.' *The Psychologist*, 23 (1) 26-27.

Link to published version:

http://www.thepsychologist.org.uk/archive/archive_home.cfm?volumeID=23&editionID=183&ArticleID=1616

Sholl, C., Korkie, J. & Harper, D. (2010). Challenging teenagers' ideas about people with mental health problems. *The Psychologist*, 23(1), 2-3.

Challenging teenagers' ideas about people with mental health problems

Catherine Sholl
Clinical Psychologist
Ealing Services for Children with Additional Needs
West London Mental Health Trust
Carmelita House
21-22 The Mall
London W5 2PJ
catherine.sholl@nhs.net

Juan Korkie
Kaya House Manager
Barnet Voice for Mental Health
Avenue House
East End Road
London N3 3QE
juankorkie@yahoo.com

Dave Harper
Reader in Clinical Psychology
School of Psychology
University of East London
Romford Road
London E15 4LZ
d.harper@uel.ac.uk

Mental health service users are one of the most socially excluded groups in society. Surveys indicate that younger people have less understanding of their own and others' mental health. In this article we discuss attempts to promote a psychosocial understanding of mental health in schools and describe a brief classroom-based intervention in a secondary school. It emphasises the importance of psychologists and service users working together to challenge discrimination against people with mental health difficulties.

Discrimination is a major barrier to the social inclusion of people with mental health problems (Social Exclusion Unit, 2004). Teenagers are an important group to target since they appear to have less tolerant attitudes about mental health (Care Service Improvement Partnership, 2009) and may even harass mental health service users (Berzins et al., 2003), something which, sadly, has figured recently in the news with Fiona Pilkington's killing of her daughter Francessca – who had a learning disability – and herself, apparently after years of harassment by local youths. Moreover, adolescence is a time when teenagers themselves may experience such problems and yet they find it more difficult to seek help than adults.

A recent longitudinal study has shown that attitudes towards people with mental health problems worsened in England – and to a lesser extent in Scotland – between 1994 and 2003 (Mehta et al., 2009). This is despite extensive campaigns (e.g. Crisp et al., 2000). One possible explanation for this is that campaigns to address discrimination against mental health service users have so far tended to be based on a biomedical model. Indeed, Angermeyer and Matschinger (2005) reported that the increased desire for social distance from people with a diagnosis of schizophrenia found in their surveys of the West German public between 1990 and 2001 was matched by a parallel increase in the tendency of the public to endorse biological causes. However, psychosocial explanations – where mental health problems are seen as caused by adverse life events – appear to be associated with more positive attitudes and behaviour (Read et al., 2006).

Compared with the research on adults, there has been relatively little investigation of young people's views of mental health. In an interesting qualitative study Lindley (2009) used photo vignettes to elicit accounts from 14- to 18-year-olds. She reported that teenagers found it easier to adopt non-discriminatory positions towards people with mental health problems when the cause of those problems was seen as lying in their life history – i.e. when a psychosocial explanatory framework was available.

Some recent studies have indicated that educational interventions based on psychosocial models of mental health may be more effective in bringing about positive attitude change. Schulze et al. (2003) developed a project week for German teenage school students during which they met a young person with a diagnosis of schizophrenia. They focused on similarities rather than differences, including the idea that life events (such as poverty, disability, violence, drugs and loneliness) were important determinants of well-being for everyone. The mental health service user discussed their experience of

mental health difficulties and the experience of discrimination resulting from having a diagnosis of schizophrenia. The project took place over five days and all of the elements were interactive with issues being explored through guided group discussions. Measures of attitudes and behaviour indicated that there were positive changes both immediately and one month after the intervention with regard to both the stereotypes the young people held and their readiness to enter social relationships with people with a diagnosis with schizophrenia.

In a similar study carried out in the UK (Pinfold et al., 2003), secondary school students attended two mental health awareness workshops. Group 1 received two hour-long sessions on understanding mental health problems from a psychosocial perspective delivered by a mental health professional. Group 2 received the same content, but with sessions co-facilitated by a mental health service user. Positive attitude scores rose significantly following psychosocial education, in particular for female students, grammar school students and those who had had personal contact with people with mental health problems.

We designed an intervention aimed at challenging mental health discrimination and delivered it to a class of 25 13-14 year old students in an inner London school (Sholl, Korkie & Harper, in press). They received four weekly sessions (each 50 minutes long) facilitated by Catherine (then a trainee clinical psychologist) and Juan (a mental health service user and clinical psychologist) – Dave's role was as Catherine's research supervisor. We took a collaborative approach whereby the group were encouraged to discuss their beliefs and reflect upon what they had learned and experienced within each session.

INSERT BOX 1 ABOUT HERE

Our approach was underpinned by two assumptions:

- The active participation of service users is important in challenging stereotypes (Pinfold et al., 2003).

- Advocating a bio-medical model is ineffective and may even maintain or increase negative attitudes towards people with mental health problems (Read et al., 2006). Instead we were influenced both by a psychosocial approach (Read et al., 2006) and by the disability inclusion model (Sayce, 2000). Here there is an emphasis on challenging discrimination in society and increasing empathy by viewing mental health problems as understandable responses to difficult life events.

Feedback from the young people involved

Quantitative and qualitative analysis indicated that, following the intervention, the young people had more positive attitudes about mental health. Key factors facilitating change included:

- A collaborative and respectful approach by the facilitators.
- Meeting a service user and hearing their story, not only in relation to mental health but also learning about them as a person (e.g. their work and hobbies).
- Receiving accurate information.

INSERT BOX 2 ABOUT HERE

Conclusion and implications

Despite the rather negative picture painted by attitude surveys, it is clear that, in the right conditions, teenagers are open to changing their views about mental health problems. This project emphasises the need for effective teaching about mental health problems in schools. Such teaching would be an innovative way of meeting some of the aims of *Every Child Matters* like

encouraging children to 'be healthy' and 'make a positive contribution'. Working alongside service users, psychologists could begin more proactive work aimed at reducing the mental health discrimination and inequality found in society (e.g. schools, employers etc). What is needed now is more extensive provision of this kind of intervention, properly evaluated, across UK schools. This would help us to develop our knowledge about effective anti-discrimination interventions. However, in order for this to occur, it is essential that there is better communication and more joint work not only between health and education agencies but also involving mental health service users. Only then will we begin to see consistent progress in both addressing mental health discrimination and improving young people's well-being.

Note: Further information on the project is available on request from the authors.

References

Angermeyer, M.C. & Matschinger, H. (2005). Causal beliefs and attitudes to people with schizophrenia. *British Journal of Psychiatry*, 186, 331–334.

Berzins, K. Petch, A. & Atkinson, J.M. (2003). Prevalence and experience of harassment of people with mental health problems living in the community. *British Journal of Psychiatry*, 183, 526-533.

Care Service Improvement Partnership (2009). *Attitudes to mental illness 2009: Research report*. London: Department of Health.

Crisp, A.H., Gelder, M.G., Rix, S. et al. (2000). Stigmatisation of people with mental illnesses. *British Journal of Psychiatry*, 177, 4–7.

Lindley, E. (2009). Gateways to mental illness discourse: Tools for talking with teenagers. *International Journal of Mental Health Promotion*, 11, 15–22.

Mehta, N., Kassam, A., Leese, M. et al. (2009). Public attitudes towards people with mental illness in England and Scotland, 1994–2003. *British Journal of Psychiatry*, 194, 278–284.

Office for National Statistics (2007). *Attitudes to Mental Illness 2007*. London: Author.

Pinfold, V. Toulmin, H. Thornicroft, G. Huxley, P. Farmer, P. & Graham, T. (2003). Reducing psychiatric stigma and discrimination: Evaluation of educational intervention in UK secondary schools. *British Journal of Psychiatry*, 182, 342-346.

Read, J. Haslam, N. Sayce, L. & Davies, E. (2006). Prejudice and schizophrenia: A review of the 'mental illness is an illness like any other' approach. *Acta Psychiatrica Scandinavica*, 114, 303-318.

Sayce, L. (2000). *From Psychiatric Patient to Citizen*. New York: St Martin's Press.

Schulze, B., Richter-Werling, M. Matschinger, H. & Angermeyer, M.C. (2003). Crazy? So what! Effects of a school project on students' attitudes towards people with schizophrenia. *Acta Psychiatrica Scandinavica*, 107, 142-150.

Sholl, C., Korkie, J. & Harper, D. (2009). Working with young people to challenge discrimination against mental health service users: A psychosocial pilot study. *Clinical Psychology Forum*, 196, 45-49.

Social Exclusion Unit (2004). *Mental Health and Social Exclusion*. London: Office of the Deputy Prime Minister.

BOX 1

Content of teaching sessions

Session 1: An open, interactive (rather than didactic) discussion about the group's ideas about people with mental health problems.

Session 2: The group interview Juan about his life and experiences of distress.

Session 3: Understanding mental health distress – exploring what we need to live and be happy, and how we might respond if these things were not available.

Session 4: Presenting information to help challenge any myths which have been raised (e.g. that all people with mental health problems are dangerous).

Box 2

Juan's reflections on service-user involvement

- Service users can often be made to feel that their involvement is tokenistic and that they are not seen as equals. However, this was avoided here because both Catherine and I were equally involved in the planning and negotiation of our respective roles.
- My dual perspective both as a service user and as a psychologist seemed to be helpful both in drawing out the students' stereotypes and in helping them to reflect on this process.
- We aimed to help the young people to move from sympathy (which implies a level of social distance) to empathy.
- Stereotypes about people experiencing distress involve several layers of socialized reactions. Therefore part of the work was to help the young people to develop their general social skills – for example being open to difference.
- It was important for the teenagers to hear about other parts of my life and so realise that being a service user is not the sum total of my identity.
- Obviously it was not possible for me to 'represent' the experience of all users of mental health services but I aimed to enable the young people to draw wider lessons from both my own unique experiences of distress and the ways in which services have responded to it.