



ORAL PRESENTATION

Open Access

# Reliability of three foot models to examine paediatric gait

Ryan Mahaffey\*, Stewart Morrison, Wendy Drechsler, Mary Cramp

From 3rd Congress of the International Foot and Ankle Biomechanics Community Sydney, Australia. 11-13 April 2012

## Background

A variety of multi-segmental foot models have been produced to examine patterns of foot segmental movement during gait cycle to identify biomechanical differences between normal and pathological foot function[1-3]. The reliability of foot models to accurately describe motion of the foot joints is dependent on the ability of the examiner to repeatedly apply markers to specific landmarks and the relevance of models' segmental descriptions to underlying anatomy. The aim of this study was to test the reliability of segmental angles measured by three published foot models during paediatric gait.

## Materials and methods

Sixteen children, aged 6 to 12 years old, were recruited to the study. Marker sets for three foot models 3DFoot [1], Oxford Foot Model (OFM)[2], and Kinfoot[3] were applied to their right feet simultaneously which to the authors knowledge, is the first direct comparison of the three models during gait. Each foot model was assessed for repeatability of maximal joint angle and range of motion during the gait cycle between two testing occasions. Absolute angular differences and standard error of measurement (SEM) are reported.

## Results

Repeatability of all maximal segmental angles and range of motions were higher in 3DFoot compared to OFM and Kinfoot (Table 1).

## Conclusion

Decreased measurement error observed in 3DFoot and Kinfoot models may be attributable to normalisation of kinematics data to subject standing position. In the

**Table 1 Inter-session repeatability of foot model's 3D maximal segmental angles over the gait cycle.**

Model	Segments	Maximal joint angle		Range of joint angle	
		Difference °	SEM °	Difference °	SEM °
OFM	Hindfoot to Shank	2.1 ± 15.1	10.9	1.2 ± 8.0	5.7
3DFoot	Hindfoot to Shank	1.0 ± 5.2	3.6	1.0 ± 4.6	3.3
Kinfoot	Hindfoot to Shank	1.0 ± 5.1	3.6	1.4 ± 6.3	4.3
3DFoot	Midfoot to Hindfoot	0.8 ± 3.5	2.2	0.3 ± 2.7	1.9
Kinfoot	Midfoot to Hindfoot	3.0 ± 11.1	6.7	3.7 ± 11.3	6.6
OFM	Metatarsals to Hindfoot	0.8 ± 8.5	5.3	1.3 ± 5.7	5.4
3DFoot	Metatarsals to Midfoot	0.7 ± 4.0	2.9	0.6 ± 3.6	2.5
Kinfoot	Metatarsals to Midfoot	2.8 ± 7.8	4.8	2.6 ± 6.6	3.7
OFM	Hallux to Metatarsals	2.3 ± 15.6	11.2	0.4 ± 13.7	9.1
3DFoot	Hallux to Metatarsals	1.5 ± 10.0	6.2	0.4 ± 12.6	8.8
Kinfoot	Hallux to Metatarsals	4.4 ± 21.8	15.1	2.1 ± 11.8	7.2

OFM, non-normalisation of gait data resulted in variable segmental offsets, particularly in the frontal plane. Greater measurement error was observed for several foot segments in the Kinfoot model. This may be due to discrepancies in model segment definitions in relation to the underlying joint anatomy, especially around the midfoot to hindfoot segments. 3DFoot model consistently showed the least measurement error in the segment motions examined and thus is appropriate for use to examine foot biomechanics in gait.

Published: 10 April 2012

## References

1. Leardini A, Benedetti M, Berti L, Bettinelli D, Nativio R, Giannini S: **Rear-foot, mid-foot and fore-foot motion during the stance phase of gait.** *Gait Posture* 2007, **25**:453-462.

\* Correspondence: r.mahaffey@uel.ac.uk  
School of Health, Sport & Bioscience, University of East London, UK

2. Carson M, Harrington M, Thompson N, O'Connor J, Theologis T: **Kinematic analysis of a multi-segment foot model for research and clinical applications: a repeatability analysis.** *J Biomech* 2001, **34**:1299-2307.
3. MacWilliams B, Cowley M, Nicholson D: **Foot kinematics and kinetics during adolescent gait.** *Gait Posture* 2003, **17**:214-224.

doi:10.1186/1757-1146-5-S1-O18

**Cite this article as:** Mahaffey *et al.*: Reliability of three foot models to examine paediatric gait. *Journal of Foot and Ankle Research* 2012 **5**(Suppl 1):O18.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

