

# **Clustering of lifestyle risk behaviours among residents of forty deprived neighbourhoods in London: Lessons for targeting public health interventions**

**Authors:** Watts P (Dr)<sup>1\*</sup>; Buck D (Dr)<sup>2</sup>; Netuveli G (Prof)<sup>1</sup>; Renton A (Prof)<sup>1</sup>

\*Corresponding Author

<sup>1</sup>Institute for Health and Human Development, University of East London, London, E15 4LZ, United Kingdom.

<sup>2</sup>The King's Fund, London, W1G OAN, United Kingdom.

## **Abstract**

**Background:** Clustering of lifestyle risk behaviours is very important in predicting premature mortality. Understanding the extent to which risk behaviours are clustered in deprived communities is vital to most effectively target public health interventions.

**Methods:** We examined co-occurrence and associations between risk behaviours (smoking, alcohol consumption, poor diet, low physical activity and high sedentary time) reported by adults living in deprived London neighbourhoods. Associations between sociodemographic characteristics and clustered risk behaviours were examined. Latent class analysis was used to identify underlying clustering of behaviours.

**Results:** Over 90% of respondents reported at least one risk behaviour. Reporting specific risk behaviours predicted reporting of further risk behaviours. Latent class analyses revealed four underlying classes. Membership of a maximal risk behaviour class was more likely for young, white males who were unable to work.

**Conclusions:** Compared to recent national level analysis there was a weaker relationship between education and clustering of behaviours and a very high prevalence of clustering of risk behaviours in those unable to work. Young, white men, who report difficulty managing on income were at high risk of reporting multiple risk behaviours. These groups may be an important target for interventions to reduce premature mortality caused by multiple risk behaviours.

## **Introduction**

Clustering of lifestyle risk behaviours is critical to health. A longitudinal study in Norfolk, UK (1) which followed 20,244 men and women for an average of 11 years found that those who had all four lifestyle risk behaviours (smoking and non-adherence to government guidelines on alcohol consumption, diet and physical activity respectively) were four times more likely to have died at follow up compared to those who had none of the lifestyle risk behaviours. Similar patterns of mortality risk for clustered behaviours have been reported in a study of 11 European countries (2) and in US cohorts (3, 4). We therefore need to urgently know the extent of clustering in the English population, and think hard about whether policies focused on behaviour change in isolation from one another, either national or local, are really going to work.

A recent Kings Fund analysis (5) of Health Survey for England data on lifestyle risk behaviours has thrown light on the distribution of lifestyle health risks in the English population, how they cluster in different populations and how this has changed over time. The core findings of the King's Fund study (5) are first that risk factors do cluster systematically by age, socioeconomic class, education and to some extent gender. Furthermore, a study of adults in the east of England has demonstrated that clustering of risk behaviors is more prevalent in deprived areas (6).

The King's Fund study (5) argued that in the new public health system both central and local government need to be much smarter in their understanding of clustering, and the policy and practice response to it if the government's stated aim of "improving the health of the poorest fastest"(7) is to be met. It also argued that local areas could make much more effective use of their own health and lifestyle surveys by applying "a clustering lens" to how lifestyles are distributed to understand the nature of clustering in local areas and what this implies for local policies and practice. This paper takes on that challenge for deprived neighbourhoods in London, using data from a 2008 household survey of more than four thousand residents in 40 of London's most deprived neighbourhoods.

## **Material and Methods**

### ***Data Source***

Data were taken from the adult baseline survey of the *Well London* cluster randomised trial (CRT), details and results of which are reported elsewhere (8-10). Within the *Well London* CRT, a pair of census lower super output areas (LSOAs) amongst the 11% most deprived (based on the English Indices of Multiple Deprivation) across London were selected in each of 20 London boroughs. This study used data from the household adult survey collected in all of the 40 selected LSOAs during 2008.

### ***Defining risk behaviours***

The Well London baseline survey dataset contains information about five unhealthy behaviours: a) Smoking status; b) Alcohol consumption; c) Low fruit/vegetable consumption; d) Low levels of physical activity e) Sedentary time. Current smoking behaviour was ascertained by a simple yes/no question “are you a daily smoker?”, as commonly used in the Health Survey for England (11). Self-reported alcohol consumption was recorded using an item from previous studies of health behaviours in London (12), dichotomised as 1=‘drink heavily’, ‘drink quite a lot’ or ‘drink a moderate amount’; 0=‘drink a little’, ‘hardly drink at all’ or ‘never drink alcohol’. Validated measures of healthy eating (food frequency questionnaire adapted from the Health Survey for England 2008 (11)) were used to create a binary indicator for meeting the Department of Health’s recommended minimum of five portions of fruit or vegetables per day (13). The international physical activity questionnaire – short form (IPAQ-SF) (14) was used to generate a binary variable analogous to meeting the levels of physical activity (5 x 30 minutes of at least moderate intensity) recommended by the chief medical officer for England (15, 16): 1=does not meet physical activity recommendations; 0=meets physical activity recommendations. Data on sedentary time was obtained using a single item from the IPAQ-SF that asks respondents to recall the total time they have spent sitting at any time on a weekday. More than four hours sitting time per day is generally considered excessive (17), therefore a binary variable was generated using IPAQ-SF data: 1= More than four hours sitting time on a weekday; 0= Four hours or less sitting time on a weekday.

### ***Data Analyses***

Data on the five risk behaviours were used to derive variables to describe the number of risk behaviours in each respondent (0-5) and patterns of risk behaviours (all possible combinations of risk behaviours). Logistic regression models were used to examine associations between risk behaviours each sociodemographic factor. Ordered logistic regression models were used to examine associations between sociodemographic factors and number of risk behaviours (count of risk behaviours). Robust standard errors were used to account for within neighbourhood clustering of the sample. Logit regression models were used to examine relationships between risk behaviours. Odds ratios (OR) from these logit regression models were converted into predicted probabilities  $\left(\frac{OR}{1+OR}\right)$ .

We used latent class analyses to discover underlying clustering of people according to the multiple risk behaviours they adopt. We tested the data for two, three, four and five classes. While there were from two to four classes the models showed significant differences and there were no differences between solutions with four classes and five classes. Therefore, we chose the four classes. We studied the behavioural profile of each class by looking at the proportion of individuals exhibiting the different behaviours. We also used multinomial logistic regression with robust standard errors to predict the relative risk of belonging to the different classes based on sociodemographic variables. All data analyses were conducted in Stata v.11 and Mplus v.7.

### ***Missing data***

Descriptive analyses were conducted using complete cases. Analyses in other steps were conducted using multiple imputation to account for missing data. Details of how the multiple imputation models were specified have been published previously (9). Proportions of missing data are shown in Supplementary File 1.

### **Results**

Ninety three percent of men and 92% of women reported at least one risk behaviour and 70% of men and 66% of women reported two or more than two risk behaviours. Supplementary File 2 shows the proportion of men and women who reported zero to five risk behaviours.

Figure 1 shows the prevalence of all possible combinations of risk behaviours, where for example, 'S' represents daily smoking but no other risk behaviours and 'SD' represents daily smoking and moderate/high alcohol consumption but no other risk behaviours. Low fruit and vegetable consumption together with high sedentary time was the most commonly reported combination of risk behaviours for both men (9.6%) and women (11.4%). In Supplementary Files 3 and 4 these figures are presented using only the four risk behaviours included in the King's Fund analysis (without the addition of sedentary time).

Evidence for the extent to which risk behaviours were 'clustered' is shown in Table 1 where predicted probabilities from logit regression models show the probability of a respondent reporting a specific risk behaviour given that they had reported another specific risk behaviour or combination of risk behaviours. For example, for respondents that reported moderate/high alcohol consumption (D) the probability of them reporting being a daily smoker (S) was 0.8. There are many instances shown in Table 1 where combinations of risk behaviours predicted other risk behaviours, for example, respondents who reported daily smoking and high sedentary time, the probability of reporting moderate/high alcohol consumption was 0.79.

Table 2 shows the association between each risk behaviour and sociodemographic factors (logistic regression) and associations between the count of risk behaviours and sociodemographic factors (ordered logistic regression). These results show that there was considerable variation in odds ratios (and thus prevalence of different individual and multiple risk factors) across groups. In terms of individual risk factors for example, women were more likely to have low physical activity, but less likely to report smoking or moderate/high alcohol consumption than men; when compared to 16-24 year olds, 25-54 year olds were more likely to report smoking and moderate/high alcohol consumption and increasing age was associated with lower physical activity levels; those unable to work reported higher prevalences of low fruit and vegetable consumption, low physical activity, high sedentary time and were more likely to smoke than those in paid employment; and rates of moderate/high alcohol consumption were uniformly lower for all ethnic groups compared to the white ethnic group.

Results from the ordered logistic regression of the count of risk behaviours showed that women had significantly lower odds of reporting a higher number of risk behaviours than

men. Compared to respondents aged 16-24 years old, those aged 35-44, 55-64 and over 65 years old had higher odds of reporting more risk behaviours. White respondents had significantly higher odds of reporting more risk behaviours when compared to all other ethnic groups. Respondents who reported being unable to work, ill or disabled had more than three times higher odds of reporting more risk behaviours compared to those working more than 30 hours per week. We did not find evidence of associations between education status or ease of managing on income and total number of risk behaviours reported.

The latent class analysis revealed four latent classes. Figure 2 shows the profile of behaviours in each class. In Class 1 there were a minimal amount (<20%) of smoking, poor diet and sedentary behaviours (class label: 'minimal behaviours'). In Class 2, the majority smoked, reported low fruit and vegetable consumption and about a third also reported high alcohol consumption (class label: 'smokers'). In the third class, all behaviours except poor physical activity were exhibited by the majority (class label: 'maximal behaviours'). Finally, in the largest class, Class 4, the majority reported high sedentary time, low fruit and vegetable consumption and low physical activity levels (class label: 'sedentary lifestyle'). The classes were labelled according to these profiles.

The sociodemographic profiles of these classes are shown in Table 2. Table 2 also shows the relative risk of belonging in each class according to sociodemographic factors. When compared to the minimal behaviours class, respondents who were younger, male, white and unable to work were at greater risk of membership of the maximal behaviours class. Being white and unable to work was associated with greater risk of membership of the 'smokers' class.

## **Discussion**

### ***Main findings of this study***

Our analyses suggest that there is clustering of risk behaviours in this survey sample in deprived London neighbourhoods. Daily smoking and moderate/high alcohol consumption were most strongly correlated and respondents reporting combinations of two, three or four risk behaviours were more likely to report further risk behaviours. Four classes of behaviours were identified using latent class analysis; 'sedentary lifestyle', 'minimal risk behaviours',

‘maximal risk behaviours’ and ‘smokers’. The largest class was the sedentary class, and those at highest risk of membership of this class were female and retired or unable to work. When compared to the minimal behaviours class, membership of the maximal risk behaviours class was associated with being younger, white, male, and unable to work.

### ***What is already known on this topic***

Our findings suggest that only 8% of respondents to the survey had none of the five lifestyle risk behaviours examined here. This is slightly higher than the King’s Fund analysis of Health Survey for England Data (5), which found 4% in 2003 and 6% in 2008. The King’s Fund analyses focussed on clustering of four behaviours, whereas our analyses have additionally included analysis of sedentary time. Although, our results and the national results from the King’s Fund study are not directly comparable for this and other reasons, our findings are worth comparing and discussing further.

We find that rates of low reported physical activity are considerably lower than in the Health Survey for England data which the King’s Fund used. The King’s Fund found a complex relationship between risk behaviours and age, and speculated that this may be due to a combination of age and survivor effects. We report similar findings, with the odds of a higher number of risk behaviours highest in 35-44, 55-64 and over 65 year olds suggestive of life-course trajectories of risk. The King’s Fund’s national study did not assess ethnicity specifically. However, we find ethnicity is important, with Black, Asian and Mixed ethnicity associated with a lower number of risk behaviours. Much of this effect may be driven by their lower smoking and alcohol consumption rates reflecting differences in cultural attitudes to tobacco and alcohol.

We found that in this population living in deprived parts of London, educational attainment does not explain risk behaviour clustering after adjusting for other sociodemographic factors. This contrasts with the national findings from the King’s Fund study. It could be that the effect of education is being picked up in other variables, for instance in the employment or income variables. However, it could also be that education is less “protective” in this population living in deprived neighbourhoods, from taking on multiple risk behaviours. Evidence from a review of large datasets on determinants risk behaviours (18) suggests that large proportions of the variance by education in individual risk behaviours can be accounted for by differences in



characteristics not examined in this study, including family background, social networks, knowledge and cognitive ability.

### ***What this study adds***

One of the starkest findings in this study is that people who report unable to work have more than three times higher odds of reporting a higher number of risk behaviours than those in full-time paid employment and more two and a half times the risk of belonging to the maximal behaviours class. Sixty seven percent of those unable to work, ill or disabled reported at least three risk behaviours. The latent class analyses also revealed that after adjusting for other sociodemographic factors, those who were not in employment were more likely to report a lifestyle characterised by high sedentary time, low levels of physical activity and low fruit and vegetable consumption. This raises politically sensitive questions of cause and effect, whether their lifestyles have led them into inability to work, or inability to work has led to poor lifestyles. Clearly, there is likely to be an element of both, and getting to the root cause will be important for designing appropriate solutions.

What is clear for local authorities as they take up their new roles for health improvement in London, is that members of these behavioural risk classes are at very high risk of premature mortality due to lifestyle clustering and should be a priority for targeted action. Is there any sign of this yet? At a strategic level we would hope to see this in Health and Wellbeing Board's strategies. The Local Government Association has recently developed a summary tool (19). Using this we searched for combinations of priorities focussing on alcohol and substance abuse, healthy living, obesity in adults, smoking, health inequalities and prevention and early intervention. All boroughs, except Brent and Richmond upon Thames mentioned at least one of these as priorities, and four, Barking and Dagenham, Hillingdon, Kingston upon Thames and Merton had four or more of these six areas as priorities. Yet, looking in depth at their strategies, the concept of how their populations experience behaviours as clusters – and the response – is either absent or inadequately expressed as part of the narrative of these core strategic plans.

This does not mean there are not pockets of relevant activity in these boroughs (for instance Merton's LiveWell service (20)) or their local NHS, or indeed elsewhere, but it seems it is not part of mainstream strategy to take clustering of behaviours into account when deciding on

how to set and implement prevention and public health plans locally. We agree with the King's Fund, that if we want to improve the health of London's poorest fastest we need to understand and target behaviour change strategies and implement them in a way that populations actually experience them, rather than relying solely on blanket single behaviours approaches, one by one. There is little sign that this is happening yet. There could be a task for the new London Health Board (21), given its role to bring leadership to issues of pan-London significance. But, the London Health Board or others will need additional insight and evidence to shape their actions.

Further research is needed to identify other determinants of clustered behaviours. Theories of motivation and addiction suggest that certain psychological characteristics may explain underlying causes of behaviours (22). Furthermore, social ecological theories (23) describing the interaction between these individual psychological characteristics and social and environmental cues to behaviour may be useful in guiding future research into determinants of clustered risk behaviours and interventions to target these clustered behaviours.

### ***Limitations of this study***

The cross-sectional design of this study prevents us from making inferences about the causal directions of the associations reported. For example it is not clear whether clustered risk behaviours are adopted simultaneously or whether the adoption of one risk behaviour subsequently leads to the adoption of further risk behaviours. The high levels of physical activity reported in this sample are likely to result from problems with overestimation from the IPAQ-SF that have recently been reported in a systematic review of validation studies (24). The measure of physical activity in the Health Survey for England that the King's Fund used is a self-report recall measure similar to the IPAQ-SF. However, unlike the IPAQ-SF, there are not substantial concerns about the validity of physical activity measure used in the Health Survey for England.

It is also worth noting some strengths of the methods applied in this study. Previous studies examining clustering of risk behaviours (25) have used approaches that have either analysed co-occurrence of risk behaviours or associations between co-occurring risk behaviours. We have used these approaches, but have additionally examined underlying clustering of behaviours through analysis of latent class membership and looked at risk of class

membership according to sociodemographic factors. This may be an additional helpful way to target activity.

**Funding**

This work was supported by the Wellcome Trust [083679/Z/07/Z]

**Acknowledgements**

We are very grateful for the contributions of numerous fieldworkers to the collection of the household survey data. We would also like to thank members of the Well London evaluation team for their contributions to the data collection and preparation processes.

**Table 1. Predicted probabilities for associations between risk behaviours.**


Existing behaviour	Smoking (S)	Drinking (D)	Low fruit/vegetable consumption (F)	Low physical activity (P)	High sitting time (T)
S					
D	<b>0.8</b>				
F	0.57	0.57			
P	0.48	0.47	<b>0.62</b>		
T	0.49	0.54	0.54	<b>0.67</b>	
SD			0.58	0.48	0.57
SF		<b>0.77</b>		0.49	0.49
SP		<b>0.72</b>	0.62		<b>0.66</b>
ST		<b>0.79</b>	0.57	0.55	
DF	<b>0.77</b>			0.54	0.52
DP	<b>0.77</b>		<b>0.67</b>		0.64
DT	<b>0.77</b>		0.55	0.51	
FP	0.48	0.51			<b>0.65</b>
FT	0.53	0.55		<b>0.66</b>	
PT	0.49	0.47	<b>0.61</b>		
SDF				0.54	0.53
SDP			0.54		0.65
SDT			0.55	0.52	
SFP		<b>0.76</b>			<b>0.68</b>
SFT		<b>0.77</b>		0.57	
SPT		<b>0.71</b>	0.66		
DFP	<b>0.74</b>				0.62
DFT	<b>0.76</b>			0.58	
DPT	<b>0.74</b>		0.65		
FPT	0.5	0.5			
SDFP					0.66
SDFT				0.58	
SDPT			0.68		
SFPT		<b>0.75</b>			
DFPT	<b>0.75</b>				

Emboldened figures represent statistically significant probabilities at P-values  $\leq 0.001$ .


**Table 2. Relationships between sociodemographic factors and risk behaviours and latent classes of risk behaviours**

Sociodemographic factor	Smoker OR	Moderate to high alcohol OR	Low fruit and veg OR	Low physical activity OR	High sitting time OR	Count of risks (ordered logistic) OR	Latent					
							Minimal behaviours			Smokers		
							N	%	RRR	N	%	
<b>Sex (Male)</b>	Ref	Ref	Ref	Ref	Ref	Ref	486	43.4	Ref	124	48.3	
Female	0.54***	0.42***	0.98	1.66***	0.90	0.76***	634	56.6	Ref	133	51.8	
<b>Age (16-24)</b>	Ref	Ref	Ref	Ref	Ref	Ref	240	23.4	Ref	44	19.9	
25-34	1.36*	1.56**	0.80*	1.24	1.06	1.16	274	26.7	Ref	79	35.6	
35-44	1.41*	1.66**	0.91	1.59***	0.93	1.37*	219	21.3	Ref	42	18.9	
45-54	1.57**	1.58*	0.68**	1.60**	0.91	1.17	132	12.9	Ref	30	13.5	
55-64	1.19	1.42	0.88	2.46***	0.93	1.37*	83	8.1	Ref	16	7.2	
65+	0.76	0.91	1.00	3.54***	1.50*	1.60***	80	7.8	Ref	11	5.0	
<b>Ethnicity (White)</b>	Ref	Ref	Ref	Ref	Ref	Ref	426	37.9	Ref	138	55.0	
Black	0.35***	0.44***	1.16	0.99	0.81	0.54***	387	34.5	Ref	53	21.2	
Asian	0.26***	0.17***	0.89	1.06	0.92	0.43***	187	16.7	Ref	24	9.6	
Mixed	0.69	0.48**	1.01	0.67	0.59**	0.52***	65	5.8	Ref	20	7.9	
Other	0.74	0.52*	0.59**	1.09	0.62	0.46***	58	5.16	Ref	16	6.4	
<b>Highest Education Level (Primary)</b>	Ref	Ref	Ref	Ref	Ref	Ref	98	9.3	Ref	24	10.0	
Secondary (GCSE or equivalent)	1.09	1.36	1.57**	0.62**	1.09	1.16	314	29.7	Ref	97	40.3	
A' Level or equivalent	0.87	1.41	1.36*	0.42***	1.37	0.96	298	28.2	Ref	60	24.9	
Higher (university degree)	0.67**	1.58*	1.07	0.42***	1.50*	0.93	335	31.7	Ref	59	24.5	
Other	0.84	0.47	0.77	0.82	0.83	0.79	11	1.0	Ref	1	0.4	
<b>Job Status (Working 30+ hours/week)</b>	Ref	Ref	Ref	Ref	Ref	Ref	344	31.9	Ref	81	33.5	
Working under 30 hours per week	0.84	0.74**	0.87	1.04	0.99	0.84	163	15.1	Ref	32	13.2	
Unpaid housework	0.96	0.41***	0.75	1.61**	0.52*	0.58**	67	6.2	Ref	22	9.1	
Full time education	0.36***	0.22***	1.23	1.05	1.04	0.61**	172	15.9	Ref	19	7.9	
Unemployed	1.31	0.57***	1.43**	1.35	0.95	1.10	194	18.0	Ref	67	27.7	
Retired	0.62**	0.49***	1.14	3.10***	1.55*	1.26	88	8.2	Ref	11	4.5	
Unable to work / ill / disabled	1.57*	0.76	1.76**	6.03***	2.67***	3.20***	22	2.0	Ref	2	0.8	
<b>Managing on income (difficult)</b>	Ref	Ref	Ref	Ref	Ref	Ref	528	49.2	Ref	149	61.1	
Easy or very easy	0.70***	1.12	0.96	0.94	1.42	1.04	546	50.8	Ref	95	38.9	

OR = Odds Ratio; RRR =




**Figure 1**



**Figure 2**


**Supplementary File 1. Proportions of missing data on risk behaviours in the Well London baseline survey.**

Variable	Observations	Missing	Total	% missing
Smoking status	4,043	64	4,107	1.6
Alcohol consumption	4,019	88	4,107	2.1
Low fruit/vegetable consumption	3,836	271	4,107	6.6
Low levels of physical activity	3,529	578	4,107	14.1




**Supplementary File 2**





**Supplementary File 3**



**Supplementary File 4**

## References

1. Khaw K-T, Wareham N, Bingham S, *et al.* Combined impact of health behaviours and mortality in men and women: the EPIC-Norfolk prospective population study. *PLoS Medicine* 2008;**5**(1):e12.
2. Knuops KT, de Groot LC, Kromhout D, *et al.* Mediterranean diet, lifestyle factors, and 10-year mortality in elderly European men and women: the HALE project. *JAMA* 2004;**292**(12):1433-9.
3. Ford ES, Bergmann MM, Boeing H, *et al.* Healthy lifestyle behaviors and all-cause mortality among adults in the United States. *Prev Med* 2012;**55**(1):23-7.
4. van Dam RM, Li T, Spiegelman D, *et al.* Combined impact of lifestyle factors on mortality: prospective cohort study in US women. *BMJ* 2008;**337**(7672).
5. Buck D, Frosini F. Clustering of unhealthy behaviours over time. Implications for policy and practise. *The King's Fund*, 2012.
6. Lakshman R, McConville A, How S, *et al.* Association between area-level socioeconomic deprivation and a cluster of behavioural risk factors: cross-sectional, population-based study. *Journal of Public Health* 2011;**33**(2):234-45.
7. Department of Health. *Healthy Lives, Healthy People: our strategy for public health in England*. Department of Health, 2010.
8. Wall M, Hayes R, Moore D, *et al.* Evaluation of community level interventions to address social and structural determinants of health: a cluster randomised controlled trial. *BMC Public Health* 2009;**9**(1):207.
9. Phillips G, Renton A, Moore DG, *et al.* The Well London program - a cluster randomized trial of community engagement for improving health behaviors and mental wellbeing: baseline survey results. *Trials* 2012;**13**(1):105.
10. Phillips G, Bottomley C, Schmidt E, *et al.* Well London Phase - 1: results among adults of a cluster-randomised trial of a community engagement approach to improving health behaviours and mental well-being in deprived inner-city neighbourhoods. *J Epidemiol Community Health* 2014; **68**(7):606-14
11. NHS Information Centre. *Health Survey for England 2008: Statistics on obesity, physical activity and diet*. NHS Information Centre, 2008.
12. Clark C, Haines MM, Head J, *et al.* Psychological symptoms and physical health and health behaviours in adolescents: a prospective 2-year study in East London. *Addiction* 2007;**102**(1):126-35.
13. Department of Health. 5 a day. 2014. (<http://www.nhs.uk/Livewell/5ADAY/Pages/5ADAYhome.aspx>). (Accessed 14/03/14 2014).
14. Craig CL, Marshall AL, Sjöström M, *et al.* International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 2003;**35**(8):1381.
15. Department of Health. *Start Active, Stay Active: A report on physical activity for health from the four home countries' Chief Medical Officers*. Department of Health, 2011.
16. Donaldson L. *Physical activity health improvement and prevention: at least 5 a week – a report from the Chief Medical Officer*. HM Stationery Office, 2004.
17. Biddle S, Cavil N, Eklund U, *et al.* *Sedentary Behaviour and Obesity: Review of the Current Scientific Evidence*. Department of Health, 2010.
18. Cutler DM, Lleras-Muney A. Understanding differences in health behaviors by education. *J Health Econ* 2010;**29**(1):1-28.

19. Local Government Association. *Health and Wellbeing Board priorities across England*. Local Government Association, 2014. ([http://www.local.gov.uk/health-and-wellbeing-boards/-/journal\\_content/56/10180/6111055/ARTICLE](http://www.local.gov.uk/health-and-wellbeing-boards/-/journal_content/56/10180/6111055/ARTICLE)). (Accessed 15/06/14).
20. LiveWell Merton, 2014 (<https://www.live-well.org.uk/merton>). (Accessed 15/06/14).
21. London Health Board. *What is the London Health Board?* . (<http://www.londonhealthboard.org.uk/about/default.htm>). (Accessed 15/06/14).
22. Walters ST, Rotgers F. *Treating substance abuse: Theory and technique*. Guilford Press, 2011.
23. Sallis JF, Owen N, Fisher EB. *Ecological models of health behavior*. In Glanz K, Rimer, B and Viswanath, K (Eds) *Health Behaviour and Health Education: Theory, Research and Practice*. Jossey-Bass, 2007.
24. Lee PH, Macfarlane DJ, Lam T, *et al*. Validity of the International Physical Activity Questionnaire Short Form (IPAQ-SF): A systematic review. *Int J Behav Nutr Phys Act* 2011;**8**(1):115.
25. McAloney K, Graham H, Law C, *et al*. A scoping review of statistical approaches to the analysis of multiple health-related behaviours. *Prev Med* 2013;**56**(6):365-71.